

The Tentacles of Small Intestinal Bacterial Overgrowth A Clinical Perspective

DR. GAETANO MORELLO
BC WOMEN'S HOSPITAL

Dr. Gaetano Morello

- Bachelor of Science (Cell Biology),
University of British Columbia, Vancouver,
BC
- Doctorate in Naturopathic Medicine (ND),
Bastyr University, Seattle, WA
- Clinical Practice, West Vancouver
- Clinician, Woman's Hospital, Vancouver,
BC, Chronic Complex Disease Clinic
- Author and Lecturer



Presentation

- ✓ Background Work
- ✓ My work with Central Sensitivity Syndromes (CSS)
- ✓ SIBO defined
- ✓ The Tentacles of SIBO
- ✓ Theorized Etiologies and links to SIBO
- ✓ Getting a “birds eye view.”

- ✓ Integrative Clinical Treatment Paths
 - i. Antibiotic Therapy
 - ii. Herbal Therapies
 - iii. Prokinetics and probiotics
 - iv. Elemental diet

My Work with CSS

- Spent 30 years as a naturopathic physician and the last 10 years treating complex chronic diseases at BC Women's Hospital here in Vancouver.
- In treating thousands of these patients, the link between gut health/SIBO is a key factor in treatment.
- This presentation is about that experience and my clinical observations.

My work with CSS

Complex Chronic Diseases

ME/CFS

FM

Chronic
Lyme


MS

MCS

Related Symptoms

1. Fatigue
2. Post exertion malaise
3. Cognitive Dysfunction
(concentration, brain fog, memory)
4. Pain
5. IBS- bloating, gas, constipation/diarrhea, GERD, **SIBO (underlying cause)**
6. Sleep (unrefreshing)

Central Sensitization Syndromes



My Work with Complex Chronic Disease Patients

Mitochondrial
Redox and
Upregulation

Restoring Gut
Health
SIBO

Sympathetic
Tone
downregulation

Centralized Pain
Restoration


SIBO

Small Intestinal Bacterial Overgrowth


- Definition:
 - Disruption of the normal small bowel bacterial population (usually number); may result in gas, bloating, flatulence, altered bowel function, malabsorption, pain, diarrhea/constipation
 - Accepted definition is when jejunal aspirate is $>10^5$ CFU/ml
- Wide array of effects
 - Direct injury, changes in function/sensation, gut immunology, permeability, and loss of brush border enzymes
- This imbalance can not only cause difficult to deal with GI issues but also lead to systemic complications.
 - Leaky gut, central sensitivity pain, skin , cognitive dysfunction (brain fog, memory), arthritis , fatigue, anxiety, depression, GERD

SIBO

Small Intestinal Bacterial Overgrowth



Tentacles of SIBO



- Something we've referenced for decades
- *What goes on in the "gut", all its complexities, has far reaching affects into the body.*
- We are at the tip of the iceberg in understanding the microbiome, and SIBO seems to be the **tangible area of treatment.**


Conditions associated with SIBO

(known tentacles)

✓ Irritable Bowel Syndrome (IBS)	✓ Acne rosacea
✓ Acid reflux	✓ Hyperthyroidism
✓ Coeliac disease	✓ Scleroderma
✓ Chronic Fatigue Syndrome	✓ Chronic Prostatitis
✓ Fibromyalgia	✓ Chronic Lyme
✓ Chronic constipation	✓ Myalgic Encephalomyelitis
✓ Inflammatory Bowel Disease (e.g. Crohn's and ulcerative colitis)	✓ Diverticulitis
✓ Restless leg syndrome	✓ Diabetes

Tentacles of SIBO

Clinical Observation



Background on Theorized Etiologies of Tentacles of SIBO

Gut epithelium and control of antigen trafficking

Oxidation Overload in Mitochondria due to PAMPS and DAMPS


LPS's and autoimmune diseases

Interference with nutrient absorption

Serotonin production barriers

Bacterial endotoxins

Supporting Theorized Etiologies



Correspondence | Published: 03 March 2016

Neuroinflammation in fibromyalgia and CRPS is multifactorial

Alex Vasquez

Nature Reviews Rheumatology 12, 242 (2016) | Download Citations

In his Review article (Neurogenic neuroinflammation in fibromyalgia and complex regional pain syndrome. *Nat. Rev. Rheumatol.* 11, 639–648; 2015)¹, Geoffrey Littlejohn ascribes neuroinflammation to a “neurogenic” origin, presumably triggered by pain and stress. However, attribution of neuroinflammation and central sensitization to a primary neurogenic origin is premature without integrating the well-documented coexistence of small intestine bacterial overgrowth (SIBO, one type of gastrointestinal dysbiosis), vitamin D deficiency, and mitochondrial dysfunction.

Littlejohn¹ notes that chronic pain has been associated with lipopolysaccharide (LPS)-stimulated proinflammatory cytokines (particularly IFN- γ and TNF); however, he does not pursue this line of thought to connect it to relevant literature showing clear evidence of gastrointestinal dysbiosis and increased intestinal permeability in patients with fibromyalgia and complex regional pain syndrome (CRPS). The gastrointestinal tract is the most abundant source of LPS, systemic absorption of which is increased by SIBO and increased intestinal permeability. In 1999, Pimentel et al.² showed that oral administration of antibiotics led to alleviation of pain and other clinical measures of fibromyalgia. In 2004, Pimentel et al.³ showed that among 42 fibromyalgia patients, all (100%) showed laboratory evidence of SIBO, severity of which correlated positively with severity of fibromyalgia. In that same year, Wallace and Hallegua⁴ showed that eradication of SIBO with antimicrobial therapy led to clinical improvements in fibromyalgia patients in direct proportion to antimicrobial efficacy. In 2008, Goebel et al.⁵ documented that patients with fibromyalgia and CRPS have intestinal hyperpermeability; mucosal “leakiness” was highest in patients with CRPS, indicating a strong gastrointestinal component to the illness. In 2013, Reichenberger et al.⁶ showed that CRPS patients have a distinct alteration in their gastrointestinal microbiome characterized by reduced diversity and significantly increased levels of Proteobacteria. LPS from Gram-negative bacteria is powerfully proinflammatory and is known to trigger microglial activation via Toll-like receptor 4; experimental studies have shown that LPS promotes muscle mitochondrial impairment, peripheral hyperalgesia, and central sensitization⁷.

Chronic Pain associated with LPS (IFN and TNF)

Positive correlation SIBO and FM pain (antibiotics)

Reichenberger showed that CRPS (complex regional pain syndrome) have distinct alteration in microbiome (reduced diversity levels of Proteobacteria)

Supporting Theorized Etiologies

Microbiome, 2017 Apr 26;5(1):44. doi: 10.1186/s40168-017-0261-y.

Fecal metagenomic profiles in subgroups of patients with myalgic encephalomyelitis/chronic fatigue syndrome.

Nagy-Szakal D¹, Williams BL¹, Mishra N¹, Che X¹, Lee B¹, Bateman L², Klimas NG^{3,4}, Komaroff AL⁵, Levine S⁶, Montoya JG⁷, Peterson DL⁸, Ramanan D⁹, Jain K¹, Eddy ML¹, Hornig M¹, Lipkin WJ¹⁰.

⊕ Author information

Abstract

BACKGROUND: Myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) is characterized by unexplained persistent fatigue, commonly accompanied by cognitive dysfunction, sleeping disturbances, orthostatic intolerance, fever, lymphadenopathy, and irritable bowel syndrome (IBS). The extent to which the gastrointestinal microbiome and peripheral inflammation are associated with ME/CFS remains unclear. We pursued rigorous clinical characterization, fecal bacterial metagenomics, and plasma immune molecule analyses in 50 ME/CFS patients and 50 healthy controls frequency-matched for age, sex, race/ethnicity, geographic site, and season of sampling.

RESULTS: Topological analysis revealed associations between IBS co-morbidity, body mass index, fecal bacterial composition, and bacterial metabolic pathways but not plasma immune molecules. IBS co-morbidity was the strongest driving factor in the separation of topological networks based on bacterial profiles and metabolic pathways. Predictive selection models based on bacterial profiles supported findings from topological analyses indicating that ME/CFS subgroups, defined by IBS status, could be distinguished from control subjects with high predictive accuracy. Bacterial taxa predictive of ME/CFS patients with IBS were distinct from taxa associated with ME/CFS patients without IBS. Increased abundance of unclassified *Alistipes* and decreased *Faecalibacterium* emerged as the top biomarkers of ME/CFS with IBS; while increased unclassified *Bacteroides* abundance and decreased *Bacteroides vulgatus* were the top biomarkers of ME/CFS without IBS. Despite findings of differences in bacterial taxa and metabolic pathways defining ME/CFS subgroups, decreased metabolic pathways associated with unsaturated fatty acid biosynthesis and increased atrazine degradation pathways were independent of IBS co-morbidity. Increased vitamin B6 biosynthesis/salvage and pyrimidine ribonucleoside degradation were the top metabolic pathways in ME/CFS without IBS as well as in the total ME/CFS cohort. In ME/CFS subgroups, symptom severity measures including pain, fatigue, and reduced motivation were correlated with the abundance of distinct bacterial taxa and metabolic pathways.

CONCLUSIONS: Independent of IBS, ME/CFS is associated with dysbiosis and distinct bacterial metabolic disturbances that may influence disease severity. However, our findings indicate that dysbiotic features that are uniquely ME/CFS-associated may be masked by disturbances arising from the high prevalence of IBS co-morbidity in ME/CFS. These insights may enable more accurate diagnosis and lead to insights that inform the development of specific therapeutic strategies in ME/CFS subgroups.

KEYWORDS: Chronic fatigue syndrome; Irritable bowel syndrome; Metabolic pathway; Metagenomic; Microbiota-gut-brain axis; Myalgic encephalomyelitis; Topological data analysis

Supporting Theorized Etiologies

GUT PERMEABILITY – CHRONIC INFLAMMATION

Stress induces endotoxemia and increasing barrier permeability

Karin de Punder* and Leo Pruimboom

Frontiers in Immunology published: 15 May 2015

“Chronic non-communicable diseases (NCDs) are the leading causes of work absence, disability, and mortality worldwide. Most of these diseases are associated with low-grade inflammation.”

“In combination with modern life-style factors, the increase in **bacteria/bacterial toxin translocation** arising from a more **permeable intestinal** wall causes a low-grade inflammatory state. We support this hypothesis with numerous studies finding **associations with NCDs and markers of endotoxemia**, suggesting that this process plays a pivotal and perhaps even **a causal role** in the development of low-grade inflammation and its related diseases.”

GROUND ZERO OF MOST HEALTH DISORDERS

Human Genome/Microbiome interactions

Abstract:

A short while ago, the human genome and microbiome were analyzed simultaneously for the first time as a multi-omic approach. The analyses of heterogeneous population cohorts showed that **microbiome components were associated with human genome variations**. In-depth analysis of these results reveals that the majority of those relationships are between immune pathways and autoimmune disease-associated microbiome components. Thus, it can be hypothesized that autoimmunity may be associated with homeostatic disequilibrium of the human-microbiome interactome. Further analysis of human genome–human microbiome relationships in disease contexts with tailored systems biology approaches may yield insights into disease pathogenesis and prognosis.

How does SIBO factor in Fibromyalgia


Bacterial LPS- causes inflammation, mitochondrial redox impairment, leads to sensitization

Bacterial produced gases, not made by humans interfering with homeostasis


Bacterial unknown chemical compounds(tryptophanase) possibly not producing gases creating false positives and no GI symptoms.

Getting A *Birds Eye View*


Some Essential Backg



Etiology of the Tentacles of SIBO




One Cell Paradox




An Overview of Etiology

Through Gut Permeability



Loss of Mucosal Immune Homeostasis

Levels of antigen trafficking



The Microbiome “we are not alone”

- **Knowledge level is *tip of the iceberg***
 - 100 trillion bacteria
 - 500-1000 different species of bacteria
 - 60% of fecal biomass is from bacteria
- **Microbiome exerts important effects on:**
 - Structure, physiology, biochemistry, immunology, maturation of vasculature, and gene expression
 - Human genome is in a sense static, microbiome is not (23,000 genes vs. 3.3 million genes)
 - Both microbiome and mitochondria are maternally inherited

- *Microbiota* describes the actual bacteria, and *microbiome* is the bacteria AND their genes. Terms can be used interchangeably

Notes:

- Key: Diversity and richness in microbiota species = healthy individuals
- Loss in species diversity is a common finding in several disease states.


Diversity Helps in:

1. Controlling micro-organisms
2. Immune system support
3. Maintaining wholeness of intestinal mucosa
4. Maintaining barrier effect
5. Digestive function


Beneficial bacteria, what they do

- ✓ Digestion-caloric extraction
- ✓ Detoxification
- ✓ Epigenomic expression e.g., butyrate and histone deacetylase inhibition (HDAC1, HDAC3)- associated with anti-inflammatory immune phenotype including decreasing pro-inflammatory cytokines (IL-6,8, TNF-alpha and NF-kappaB).
- ✓ Immunomodulatory cell signaling
- ✓ Cytokine modulation – insulin/leptin, interleukin 10
- ✓ Vitamin modification
- ✓ Neurotransmitters
- ✓ SCFAs and gut hormones/permeability
- ✓ Talk to the vagus nerve


A Key, Short Chain Fatty Acids



Factors influencing composition and distribution of Microbiome



Bacterial distribution pH levels



Food and
the
microbiome
(Burkina
Faso)

75% of food in Western diet is of limited or no benefit to the microbiome of the lower gut.

Refined CHO's absorbed proximally

What reaches the large intestine has limitations; small amounts of the minerals, vitamins and other nutrients necessary for the maintenance of the microbiota.

Nutrients 2013. 5, 162-207; doi:
10.3390/nu5010162

Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa

Carlotta De Filippo¹, Duccio Cavalieri², Monica Di Paola³, Matteo Ramazzotti⁴, Jean Baptiste Poullet⁵, Sebastien Massart⁶, Silvia Collini⁷, Giuseppe Pieraccini⁸, and Paolo Lionetti^{9,10}

¹Department of Preclinical and Clinical Pharmacology, University of Florence, 50139 Firenze, Italy; ²Department of Pediatrics, Meyer Children Hospital, University of Florence, 50139 Firenze, Italy; ³Department of Biomedical Sciences, University of Florence, 50134 Firenze, Italy; ⁴DNA Vision Agrifood S.A., 8-4000 Liege, Belgium; and ⁵Centre Interdisciplinaire de Spéziométrie de Masse, University of Florence, 50139 Firenze, Italy

Edited* by Daniel L. Hartl, Harvard University, Cambridge, MA, and approved June 30, 2010 (received for review April 29, 2010)

Gut microbial composition depends on different dietary habits just as health depends on microbial metabolism, but the association of microbiota with different diets in human populations has not yet been shown. In this work, we compared the fecal microbiota of European children (EU) and that of children from a rural African village of Burkina Faso (BF), where the diet, high in fiber content, is similar to that of early human settlements at the time of the birth of agriculture. By using high-throughput 16S rDNA sequencing and biochemical analyses, we found significant differences in gut microbiota between the two groups. BF children showed a significant enrichment in Bacteroidetes and depletion in Firmicutes ($P < 0.001$), with a unique abundance of bacteria from the genus *Prevotella* and *Xylanibacter*, known to contain a set of bacterial genes for cellulose and xylan hydrolysis, completely lacking in the EU children. In addition, we found significantly more short-chain fatty acids ($P < 0.001$) in BF than in EU children. Also, *Enterobacteriaceae* (*Shigella* and *Escherichia*) were significantly underrepresented in BF than in EU children ($P < 0.05$). We hypothesize that gut microbiota coevolved with the polysaccharide-rich diet of BF individuals, allowing them to maximize energy intake from fibers while also protecting them from inflammations and noninfectious colonic diseases. This study investigates and compares human intestinal microbiota from children characterized by a modern western diet and a rural diet, indicating the importance of preserving this treasure of microbial diversity from ancient rural communities worldwide.

metagenomics | nutrigenomics | biodiversity | 454-pyrosequencing | short-chain fatty acids

The human gut "metagenome" is a complex consortium of trillions of microbes, whose collective genomes contain at least 100 times as many genes as our own eukaryote genome (1). This essential "organ," the microbiome, provides the host with enhanced metabolic capabilities, protection against pathogens, education of the immune system, and modulation of gastrointestinal (GI) development (2).

We do not yet completely understand how the different environments and wide range of diets that modern humans around the world experience has affected the microbial ecology of the human gut.

Contemporary human beings are genetically adapted to the environment in which their ancestors survived and which conditioned their genetic makeup. In mammals, both diet and phylogeny influence the increase in bacterial diversity from carnivore to omnivore to herbivore (3). Dietary habits are considered one of the main factors contributing to the diversity of human gut microbiota (2). Profound changes in diet and lifestyle conditions began with the so-called "Neolithic revolution" with the introduction of agriculture and animal husbandry $\approx 10,000$ y ago (4). After that time, food resources became more abundant and constant, the concentration of large populations in limited areas

created selective pressure that favored pathogens specialized in colonizing human hosts and probably produced the first wave of emerging human diseases (5). It has been hypothesized that bacteria specialized in human-associated niches, including our gut commensal flora, underwent intense transformation during the social and demographic changes that took place with the first Neolithic settlements (6).

Western developed countries successfully controlled infectious diseases during the second half of the last century, by improving sanitation and using antibiotics and vaccines. At the same time, a rise in new diseases such as allergic, autoimmune disorders, and inflammatory bowel disease (IBD) both in adults and in children has been observed (5), and it is hypothesized that improvements in hygiene together with decreased microbial exposure in childhood are considered responsible for this increase (7). The GI microbiota plays a crucial role in the pathogenesis of IBD (8), and recent studies demonstrate that obesity is associated with imbalance in the normal gut microbiota (9, 10).

The aim of this study was to compare the gut microbiota of children aged 1–6 y living in a village of rural Africa in an environment that still resembles that of Neolithic subsistence farmers with the gut microbiota of western European children of the same age, eating the diet and living in an environment typical of the developed world. These two childhood populations provided an attractive model for assessing the impact of many environmental variables on the gut microbiota.

In our study, we address three general questions regarding the geography and evolution of the human microbiota: (i) how is bacterial diversity partitioned within and between the two populations studied; (ii) is there a possible correlation between bacterial diversity and diet; and (iii) what is the distribution of well-known bacterial pathogens in the two populations, given the different hygienic and geographic conditions?

Results and Discussion

Characterization of Dietary Habits of Children from the Boulpou Rural Village and from Florence, Italy

In this study, we characterized the fecal microbiota of 14 healthy children from the Mossi ethnic

Author contributions: C.D.F., D.C., and P.L. designed research; C.D.F., M.D.P., S.M., and S.C. performed research; S.P. contributed new reagents/analytic tools; M.S. and J.B.P. analyzed data; and C.D.F., D.C., M.D.P., and P.L. wrote the paper.


The authors declare no conflict of interest.

*This Direct Submission article had a prearranged editor.

Freely available online through the PNAS open access option.


Data deposition: Data were submitted to the Sequence Read Archive (SRA) using SRA tool (Shane and Sanger), <http://sra.broadinstitute.org/netbio/104/>. The dataset is available at <http://www.ncbi.nlm.nih.gov/sra/PR000000000>.

To whom correspondence should be addressed: E-mail: paolo.lionetti@unifi.it. This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1009610107/-/DCSupplemental.



BF

- Prevotella* } Bacteroidetes
- Xylanibacter* }
- Acetivomaculum* } Firmicutes
- Faecalibacterium* }
- Subdoligranulum* }
- Others



EU } *Alistipes* } Bacteroidetes

Burkina Faso and European children






Fig. 3. SCFA-producing bacteria could help to prevent establishment of some potentially pathogenic intestinal bacteria. (A) Quantification of SCFAs in fecal samples from BF and EU populations by SPME-GC-MS. (B) Number of sequences relative to principal *Enterobacteriaceae* genera, in BF and EU children microbiota. Mean values (\pm SEM) are plotted. Asterisks indicate significant differences (one-tailed Student *t* test of all data points: * $P < 0.05$; ** $P \leq 0.01$; *** $P \leq 0.001$).

Extrinsic and Intrinsic factors Affecting the distribution and composition of the microbiome



Migrating motor complexes (MMCs).



Migrating motor complexes (MMCs). Note that the complexes move down the gastrointestinal tract at a regular rate during fasting, that they are completely inhibited by a meal, and that they resume 90-120 minutes after the meal

Migrating Motor Complex

- Begins 90 minutes after eating
- Cleansing Wave: Waves from stomach through small intestine
- During night: 3-4 waves so have clean SI when waking
- Turned OFF during eating: DO NOT GRAZE
- Eat three meals at least 4-5 hours apart

MMC
*Causes
leading to
dysfunction*

- Slowing of Migrating Motor Complex
 - DDX
 - Morphine/opiates
 - Mixed meal
 - Stress
 - Eating
 - Diabetic neuropathy
 - Ehlers-Danlos Syndrome
 - Adhesions
 - Small gut diverticula
 - Blind loops (gastric bypass patients)
 - Narcotic use
 - Tumors of bowel
 - Extra loops of small bowel
 - Small gut obstruction

SIBO: Review Clinical Presentation

Observation and symptoms are key identifiers

- Symptoms of patients usually give enough clues
- Testing can sometimes give false negatives
- No symptoms, could still have SIBO.....note...only tip of the iceberg.

Other Symptoms

- Abdominal Pain
- Bloating/Flatulence
- Diarrhea
- Malabsorption syndrome
- Increased GUT permeability
- IBS
- Ferritin?

Flo TH, Smith KD, Sato S, Rodriguez DJ, Holmes MA, et al. (2004) Lipocalin 2 mediates an innate immune response to bacterial infection by sequestering iron. *Nature* 432: 917-921.


Background
What keeps
things in
check

HCL

Bile, Enzymes

MMC

Microbiome



Immune System

Failure of these systems can contribute to the formation of SIBO

SIBO Testing

Currently 2 main useful tests: .

1. Breath: Lactulose or Glucose or both
2. Blood: Cdt B and Vinculin Antibodies

Breath:

Note: Lactulose and Glucose have different parameters for testing positive

Interpretation:

- a. Hydrogen: Positive with a rise of 20 ppm over the lowest preceding level (90-100 minutes) for lactulose.
- b. Hydrogen: Positive with a rise of 12 ppm over the lowest preceding level (90-100 minutes) for glucose
- c. Methane: Positive with a rise of 12 ppm over the lowest preceding level (90-100 minutes) for both glucose or lactulose.

Integrative Clinical Treatment Plans for the Tentacles

Sympathetic
response
downregulation

Anti-Microbial
Therapy

Prokinetic
Support

Gut Repair

Dietary
Considerations

Elemental Diet

Treatment with Antibiotics

Consists of 2 parts:

Antibiotic for 14 days

- Rifaxamin 550 mg tid
- * Add Neomycin 500 mg bid (for methane/constipation)

Followed by Prokinetic

- Erythromycin 50 mg at night for 3 months

Herbal Antibiotic Protocol

Hydrogen
Only

Berberine (500mg): 3 capsules t.i.d.
Oregano(180 mg): 2 softgels b.i.d

Methane

Berberine (500mg): 3 capsules t.i.d.
Garlic (allicin) (500 mg): 3 softgels b.i.d.

Hydrogen
and
Methane

Berberine (500mg): 3 capsules t.i.d.
Garlic (allicin)(500mg): 3 softgels b.i.d.
Oregano (180mg): 2 softgels b.i.d.

Prokinetics

Prokinetics induce activity of the MMC and help prolong remission.

Used to help prevent relapse of SIBO

Should be started immediately after finishing treatment

Pharmaceutical:

1. Erythromycin (50 mg): 1 tablet nightly for 90 days.
2. LDN (low dose naltrexone); 1.5 mg

Botanical Approach

1. Ginger: 1000 mg per day
2. 5HTP: 50 mg at night (note medications)
3. ECPO: 3 softgels at night

Enteric Coated Peppermint oil

Remains one of the most under rated and under utilized natural “gut” products

Measurable outcomes for gas/bloating on first dose.

Key is delivery of product to S.I. through enteric coating

pH of stomach important determining factor on efficacy

Test with 1 softgel

Proper dosing: 3-4 capsules at one time

Gut Permeability
PEA
(palmitoylethanolamide)

Palmitoylethanolamide (PEA), an endogenous fatty acid amide, has been demonstrated to bind to a receptor in the cell nucleus – the peroxisome proliferator-activated receptor – and performs a great variety of biological functions related to chronic and neuropathic pain and inflammation, as has been demonstrated in clinical trials.

These include peripheral neuropathies such as diabetic neuropathy, chemotherapy-induced peripheral neuropathy, carpal tunnel syndrome, sciatic pain, osteoarthritis, low-back pain, failed back surgery syndrome, dental pains, neuropathic pain in stroke and multiple sclerosis, chronic pelvic pain, postherpetic neuralgia, and vaginal pains.

PEA is an endogenous modulator

Latest research shows PEA has an impact in reducing gut permeability

A RANDOMISED DOUBLE BLIND CONTROLLED TRIAL EXAMINING THE EFFECT OF PEA AND CBD ON THE PERMEABILITY OF THE HUMAN GUT IN VIVO

DG Couch, C Ortori, D Barrett, JN Lund and SE O'Sullivan
School of Medicine, Faculty of Science, University of Nottingham





Figure 1 – The concentration ratios of urinary lactulose and mannitol over time in healthy participants treated with aspirin and either placebo, CBD or PEA, measured by LC MS. Results are expressed as mean ratios \pm SEM. Time points between groups were compared using two-way ANOVA using Dunnett's multiple comparisons test comparing to placebo at the same time point (*p

Some Supportive Nutrients



NAC	PhamaGaba	BB536	Melatonin
1.2 g/day	200 mg	1 notte	2.5-5 mg notte

Supportive Nutrients

1. NAC : Acts as a biofilm disruptor. NAC has been shown to breakdown biofilms. Although the use of Berberine can disrupt biofilms this could be an add on.
Dosage: 1200 mg per day
2. Melatonin: 2-5 mg at bedtime can improve motility and help with sleep.
3. Probiotic: 5 billion cfu BB536: 1 capsule at night (Antibody effect)
4. Vitamin B12 (1000 mcg): Take one sublingual tablet daily. SIBO patients have low Vitamin B12 due to issues with intrinsic factor.
5. Iron Factors: 1 tablet daily.
6. PharmaGABA (100 mg): Helps up regulate parasympathetic response. Start at 100 mg 2x per day, move to 200 mg 2x per day.

Elemental Diet

Background:

- Most studied strategy in dietary management
- Has been shown to be effective in reducing relapses in Crohn Disease patients

Gastroenterology, 2017 Dec;47(12):1515-1525. doi: 10.1007/s00055-017-1543-6. Epub 2017 May 22.

Adherence to an elemental diet for preventing postoperative recurrence of Crohn's disease.

Ohara N¹, Mizushima T^{2,3}, Hama J⁴, Takahashi H¹, Iijima S^{5,6,4}, Harayuchi N¹, Inoue T⁴, Nishimura J¹, Shiraki S⁴, Iida T¹, Masuda C⁷, Yamamoto H^{1,6}, Doi K¹, Mori M¹

© Author information

Abstract

PURPOSE: An elemental diet (ED) can suppress inflammation in patients with Crohn's disease (CD); however, adherence to this diet is difficult. We examined the correlation between ED adherence and the postoperative recurrence of CD.

METHODS: The subjects of this study were 38 patients who underwent intestinal resection with anastomosis. We defined ED adherence as consuming the average daily ED dose (≥800 kcal/day) for 2 years after surgery. Patients who did not adhere to the ED were allocated to the non-ED group. We diagnosed symptomatic recurrence using the CD activity index and endoscopic recurrence using the Rutgeerts' score.

RESULTS: The ED and non-ED groups comprised 21 and 17 patients, respectively, with ED adherence of 55.3% (21/38). At the initial endoscopy, symptomatic and endoscopic recurrence rates were 4.8 and 14.3%, respectively, in the ED group, and 23.5 and 41.2%, respectively, in the non-ED group (P = 0.152 and P = 0.078, respectively). The overall symptomatic recurrence-free duration was significantly longer than the endoscopic recurrence-free duration (P = 0.022). Symptomatic and endoscopic recurrence-free durations were longer in the ED group than in the non-ED group (P = 0.003 and P = 0.021, respectively), and ED adherence was a prognostic factor for endoscopic recurrence (HR = 2.777, 95% CI = 1.036-8.767, P = 0.042).

CONCLUSION: Maintaining ED adherence for 2 years after surgery improved the symptomatic and endoscopic recurrence-free durations.

KEYWORDS: Adherence; Crohn's disease; Elemental diet; Postoperative recurrence

PMID: 28534264 DOI: 10.1007/s00055-017-1543-6

(Indexed for MEDLINE)

Aliment Pharmacol Ther; 2006 Nov 1;24(9):1333-40.

Effectiveness of an 'half elemental diet' as maintenance therapy for Crohn's disease: A randomized-controlled trial.

Tokagi S¹, Utsunomiya K, Kuriyama S, Yokoyama H, Takahashi S, Iwaouchi M, Takahashi H, Takahashi S, Kinouchi Y, Hiwatashi N, Furuyama Y, Sasaki I, Tsuji J, Shimozogawa T

© Author information

Abstract

BACKGROUND: Although thiopurines have a proven role in maintenance therapy for Crohn's disease, an alternative therapy is needed for patients intolerant or resistant to thiopurines.

AIM: To evaluate the effectiveness of home enteral nutrition as a maintenance therapy regimen in which half of the daily caloric requirement is provided by an elemental diet and the remaining half by a free diet. We refer to this home enteral nutrition therapy as 'half elemental diet'.

METHODS: Between 2002 and 2005, 51 patients in remission from two hospitals were randomly assigned to a half elemental diet group (n = 26) or a free diet group (n = 25). The primary outcome measure of this study was the occurrence of relapse over the 2-year period.

RESULTS: The relapse rate in the half elemental diet group was significantly lower [34.6% vs. 64.0%; multivariate hazard ratio 0.40 (95% CI: 0.16-0.98)] than that in the free diet group after a mean follow-up of 11.9 months. Compliance was similar in the two groups. No adverse event occurred in any of the patients throughout the study.

CONCLUSION: This randomized-controlled trial shows the effectiveness of a half elemental diet, which is a promising maintenance therapy for Crohn's disease patients.

Elemental Diet

(Overview)

BACKGROUND:

- Medical Food Beverage
- Predigested, proximally absorbed
- Supports GI disorders (helps with malabsorption and gut rest)
- Literature shows use to treat IBD (Crohn's, UC), Celiac, SIBO, Pancreatitis, GI damage (Radiation enteritis)
- Use it in hospital for nutritional support:
 - Enteral Nutrition: feeding tube or oral
 - Parenteral Nutrition-IV feeding

COMPOSITION:

Macronutrients:

- Proteins – single form amino acids
- CHO's- simple sugars
- Fats- MCT's, safflower, olive
- Micronutrients: Vitamins, Minerals and electrolytes

Absent:

1. Whole Protein
2. Fiber, Gums
3. Food items supplying micronutrients

D. Elemental Diet for SIBO (key points)

1. Used as an alternative to Abx/Habx
2. Pimental (2004), demonstrated as effective as Abx
3. Starves bacteria but feeds patient
4. Consumed in place of all other foods for 2-3 weeks
5. Significantly reduces gas after 2-3 weeks (over 130 ppm) and impacts both

Dig Dis Sci. 2004 Jan;49(1):73-7.

A 14-day elemental diet is highly effective in normalizing the lactulose breath test.

Pimentel M¹, Constantino T, Kong Y, Bajwa M, Rezaei A, Park S.

Author information

Abstract

Treatment of small intestinal bacterial overgrowth is frustrated by the low efficacy of antibiotics. Elemental diets have been shown to reduce enteric flora. In this study, we evaluate the ability of an elemental diet to normalize the lactulose breath test (LBT) in IBS subjects with abnormal breath test findings. Consecutive subjects with IBS and abnormal LBT suggesting the presence of bacterial overgrowth underwent a 2-week exclusive elemental diet. The diet consisted of Vivonex Plus (Novartis Nutrition Corp., Minneapolis, MN) in a quantity based on individual caloric requirement. On day 15 (prior to solid food), subjects returned for a follow-up breath test and those with an abnormal LBT were continued on the diet for an additional 7 days. The ability of an elemental diet to normalize the LBT was determined for days 15 and 21. A chart review was then conducted to evaluate any clinical benefit 1 month later. Of the 93 subjects available for analysis, 74 (80%) had a normal LBT on day 15 of the elemental diet. When those who continued to day 21 were included, five additional patients normalized the breath test (85%). On chart review, subjects who successfully normalized their breath test had a 66.4 +/- 36.1% improvement in bowel symptoms, compared to 11.9 +/- 22.0% in those who failed to normalize (P < 0.001). An elemental diet is highly effective in normalizing abnormal LBT in IBS subjects, with a concomitant improvement in clinical symptoms.

PMID: 14992438

[Indexed for MEDLINE]

Elemental Diet Protocols

Full Elemental Diet

Full Elemental Diet: Patient consumes 100% of caloric requirements using ED. Sole source of nutrition for the designated time period 14-21 days (as evidenced by clinical trials).

Test on Day 15, if (+) result, diet can be extended for another week. If (-) begin follow up protocol

Application: Crohn's Disease, SIBO, normalizes LBT (lactose breath test) in IBS patients.

Dosage: The dosage is approximately 1800-2000 calories. Calculate using BMR and Harris-Benedict equation (see below).

Take approximately 200-300 calorie servings every 2 to 3 hours over a 30 minute period till the caloric requirements are met (helps with blood sugar regulation).

Duration: Two weeks has been clinically validated, however, if more time is needed the physician can make that determination based on LBT outcomes.

Half Elemental Diet

Half Elemental Diet: Patient consumes 50% of daily caloric needs from ED and the other 50% from whole foods. Maintain remission from Crohn's Disease. Used when compliance becomes difficult for patients on Full ED's for SIBO and IBS. Half ED's can also be used as starting and exiting conduits to Full ED's easing the patient experience and possibly improving compliance.

Application: Maintaining remission of Crohn's after completion of Full ED, used as conduits to Full ED's and in place of ED's for difficult compliant patients. Again this will be at the discretion of physician.

Dosage: The dosage is approximately 900-1000. Calculate using BMR and Harris-Benedict equation (see below) to calculate total caloric requirement (divide this by half to give you the calories needed from the Half ED).

Duration: There are no published reports regarding the duration of a Half ED, however, 4-6 weeks can be a good starting point. The duration would be calculated at the discretion of the physician taking into account various patient symptomology and other markers deemed important.

No food or beverage during ED, however in specific cases there can be continued observable therapeutics effect with the addition:
Chicken or steak (no fat), herbal or Black tea, coffee

Elemental Diet

(Follow up after completion)

Good Follow up after ED:

To prevent bloating and help with motility

Prokinetics with meals

- Ginger; 500 mg with each meal
- Eberogast: 1 ml (20 drops) three times per day with meals
- Prescription medication at night

Transition Diet

- Day 1-2: No fiber, meats, eggs, lactose free dairy
- Day 2-3 : Add cooked pureed low FODMAP/fiber veggies (carrots, zucchini)
- Day 4 : Back to Whole Foods diet

Elemental Diet Mechanisms of Action

The ED has numerous mechanisms of action imparting the benefits attained. The following factors have been proposed as possible mechanisms of action:

1. *proximal absorption* (early assimilation of pre-digested nutrients)
2. *nutritional effects* (correction of malnutrition)
3. *low residue* (resulting from proximal absorption of near monomers and the absence of fiber)
4. *bowel rest* (another potential mechanism for the ED's ability to induce remission in IBD)
5. *decreased antigenicity* (due to the absence of antigenic whole proteins, small peptides, and particles)
6. *decreased malabsorption* (possibly resulting from the ED's ability to eradicate SIBO, which can be produced by IBD)
7. *alteration of the microbiota* (possibly a central mechanism of action¹⁶)
8. *decreased intestinal permeability*
9. *decreased proinflammatory cytokine response, may be increasing levels of interleukin 10 and Nfkappa B inhibitor*

Typical Case

CCDP program BC Women's Hospital

A 40 yr old female diagnosed with ME/CFS, on polypharmacy circa 8 years, reports 5 key symptoms:

1. Fatigue with post exertion malaise
2. Cognitive dysfunction: brain fog, memory
3. Sleep dysfunction
4. GI issue: bloating/gas, constipation/diarrhea

She reports abdominal bloating and discomfort after eating various foods, abdominal cramping, feels better when she avoids foods.

Elevation in stress levels intensifies her symptoms.

Typical Case

CCDP program BC Women's Hospital

A 60 year old woman comes in with a diagnosis of ME/CFS, FM, MCS. She has suffered with symptoms since her twenties. Claims symptoms started after a bout with mononucleosis. Further investigation demonstrate gut dysfunction after trip to third world country .

Top 5 symptoms:

1. Fatigue with post exertion malaise
2. Pain
3. GI issues: bloating/gas, abdominal pain, constipation/diarrhea
4. Cognitive dysfunction
5. Sleep issues

Currently on polypharmacy. Although they help somewhat with sleep, it remains unrefreshing. Pain is debilitating but medication just don't seem to work. Stress makes everything worst. Meditation helps.

Antibiotics during concomitant conditions have helped some symptoms.

Questions